

Animate

Animal Health

FOR ANIMAL USE ONLY

COXIBAN

Reg. No. G4166 (Act 36 of 1947)

GRANULATED ANTICOCCIDIAL AND GROWTH PROMOTANT PREMIX

INDICATIONS

For the prevention of coccidiosis in broilers and replacement pullets (maximum 12 weeks of age).

A remedy to improve mass gain and feed conversion in ruminants, and prevent coccidiosis in sheep, goats, calves and pigs.

An aid for improved feed conversion and the promotion of growth in pigs.

HARMFUL

STORAGE

Store in a cool, dry place.

COMPOSITION

Salinomycin sodium 120 g/kg

WARNINGS

Handle with care. Poisonous when swallowed.

Do not feed to horses, other equines, rhinoceros or turkeys. May be Fatal.

Do not use in dairy cows.

Do not feed to pigs 3 days prior to slaughter for human consumption.

Tiamulin (Tiamutin or Dynamutin) should not be administered when **COXIBAN** premix is being fed in the ration.

Keep out of the reach of children, animals and uninformed persons.

Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

WITHDRAWAL PERIOD

Pigs Four (3) days

Broilers and replacement pullets,
Cattle, Calves, Sheep and Goats Zero (0) days

PRECAUTIONS

Operators should use protective clothing and masks when handling.

Wash hands thoroughly after use.

Medicated feed must be used within 3 months after mixing.

DIRECTIONS FOR USE

Use only as directed. Intended for further processing. To be mixed into feed before use.

Thoroughly mix the required amount of premix according to the instructions below.

DOSAGE

Broilers:

Mix 500 g **COXIBAN** in 1,000 kg of balanced feed for a concentration of 60 ppm.

Replacement pullets:

Mix 417 g **COXIBAN** in 1,000 kg balanced feed for a concentration of 50ppm up to age 12 weeks.

Sheep, Goats and Cattle:

Complete feeds

	Active ingredient: mg Salinomycin/kg complete feed	COXIBAN premix g/ton of complete feed
Cattle	20	167
Sheep / Goats	15-20	125-167

Concentrates and finishing rations

	Quantity Salinomycin into final ration after mixing (mg active/kg feed)	Quantity COXIBAN premix into final ration after mixing (g/ton)
Cattle	20	167
Sheep / Goats	15-20	125-167

Licks

	Active ingredient: mg Salinomycin/animal/day	COXIBAN premix mg/animal/day
Cattle	20	167
Sheep / Goats	15-20	125-167

Note: Actual levels in the lick must be calculated according to intake.

Pigs:

Mix 417 g **COXIBAN** premix per ton of feed for growing pigs and 750 g of **COXIBAN** premix per ton of feed in creep rations for suckling pigs. Feed whenever necessary in creep and growing rations.

BATCH NUMBER

EXPIRY DATE

NET WEIGHT: 25 kg

Registration holder: **Animate Animal Health (Pty) Ltd.** (Co.Reg.No.1973/002312/07) 43 Hornbill Avenue, Rooihuiskraal, 0157; PO Box 66698, Highveld, 0169; Phone (+27 12) 661 3485; Fax (+27 12) 661 3486; www.animate.co.za



Animate

Animal Health

VIR DIERLIKE GEBRUIK ALLEEN

COXIBAN

Reg. Nr. G4166 (Wet 36 van 1947)

GEGRANULEERD ANTIKOKSIDIESE EN GROEISTIMULANT VOORMENSGEL

INDIKASIES

Vir die voorkoming van koksidiöse in braaikuikens en vervangingshennetjies (maksimum 12 weke ouderdom).
'n Produk om gewig en voeromsetting te verbeter by herkouers, en om koksidiöse te verhoed in skape, bokke, kalwers en varke.
'n Hulpmiddel vir verbeterde voer omsetting en groei by varke.

SKADELIK

BERGING

Berg in 'n koel, droë plek.

SAMESTELLING

Salinomisien natrium 120 g/kg

WAARSKUWINGS

Hanteer versigtig. Giftig wanneer ingeneem word.
Moenie gebruik in perde, ander perd spesies, renosters en kalkoene nie. Mag dodelik wees.
Moenie gebruik in melkkoeie nie.
Moenie in varke, 3 dae voor slag vir menslike gebruik nie
Gebruik van Tiamulin (Tiamutin of Dynamutin) nie aanbeveel tydens gebruik van **COXIBAN** voormengsel nie.
Hou buite die bereik van kinders, diere en oningeligte persone.
Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

ONTTREKKINGS PERIODE

Varke	Drie (3) dae
Braaikuikens, vervangingshennetjies, Beeste, Kalwers, Skape en Bokke	Nul (0) dae

VOORSORGMATREËLS

Werkers moet beskermende klere en maskers dra tydens gebruik van die produk.
Was hande deeglik na gebruik.
Gebruik gemengde voer binne 3 maande na vermenging.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.
Vir gebruik vir verdere verwerking. Meng met voer voor gebruik.
Meng die voorgeskrewe hoeveelheid **COXIBAN** mikro gegranuleerde voormengsel soos aangedui hieronder.

DOSIS

Braaikuikens:

Meng 500 g **COXIBAN** in 1,000 kg gebalanseerde voer vir 'n konsentrasie van 60 dpm.

Vervangingshennetjies:

Meng 417 g **COXIBAN** in 1,000 kg gebalanseerde voer vir 'n konsentrasie van 50 dpm tot en met 12 weke ouderdom.

Skape, bokke en beeste.

Volledige voer		
	Aktiewe bestaandeel: mg salinomisien/kg volledige voer	COXIBAN voormengsel g/ton volledige voer
Beeste	20	167
Skape / Bokke	15-20	125-167
Konsentrate en afrondingsrantsoene		
	Hoeveelheid salinomisien in finale rantsoen na vermenging (mg aktief/kg voer)	Hoeveelheid COXIBAN voormengsel in finale rantsoen na vermenging (g/ton)
Beeste	20	167
Skape	15-20	125-167
Lekke		
	Aktiewe bestaandeel mg salinomisien/dier /dag:	COXIBAN voormengsel mg/dier/dag:
Beeste	20	167
Skape	15-20	125-167

Let wel: Werklike vlakke in die lek moet uitgewerk word volgens die inname.

Varke:

Meng 417 g **COXIBAN** voormengsel per ton voer vir groeiende varke en 750 g **COXIBAN** voormengsel per ton voer in kruip rantsoen voor speen. Voer soos nodig in kruip en groei rantsoene.

LOT NOMMER:

VERVAL DATUM:

NETTO GEWIG: 25 kg

Registrasie houer: Animate Dieregesondheid (Edms)Bpk (Co.Reg.Nr.1973/002312/07) 43 Hornbill Laan, Rooihuiskraal, 0157;
Posbus 66698, Highveld, 0169; Foon (+27 12) 661 3485; Faks (+27 12) 661 3486; www.animate.co.za

