

Animate

Animal Health

FOR ANIMAL USE ONLY

FLAVO-8

Reg. No. G4151 (Act 36 of 1947)

FEED PREMIX

INDICATIONS

A feed additive to be used exclusively as a growth stimulant, and for the improvement of the feed conversion rate in farm animals, fish and shrimp.

Assists in restoring sensitivity of multi-drug resistant *Salmonella typhimurium* and *Escherichia coli* in farm animals

CAUTION

STORAGE INSTRUCTIONS

Store in a cool, dry place.

COMPOSITION

Flavophospholipol 8 g/kg

WARNINGS

Keep out of reach of children, uninformed persons and animals.

Irritating in the event of skin contact

Store away from food and feed.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a

wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

It is advisable to wear goggles during mixing of the product. Avoid breathing of dust. Do not eat, drink or smoke after handling the product, before washing hands and face thoroughly.

WITHDRAWAL PERIOD

Meat:

Beef Cattle, pigs, sheep, goats and broiler chickens – Zero (0) days

Milk:

Dairy Cattle – Zero (0) days

Eggs:

Layer Chickens – Zero (0) days

DIRECTIONS FOR USE

Use only as directed

It is essential that the **FLAVO-8** premix be thoroughly mixed with the feed. The following rates are recommended for general use:

Balanced feed	Flavophospholipol (active ingredient) ppm	FLAVO-8 Premix g per 1,000 kg feed (8 g active per kg)
Milk replacers	8 - 16	1000 - 2000
Calves	8 - 16	1000 - 2000
Complete cattle feed	3 - 20	375 – 2500
Dairy	3 - 20	375 – 2500
Sheep and goat feed	3 - 20	375 – 2500
Piglets	8 - 20	1000 – 2500
All other Pig diets	3 - 20	375 – 2500
Laying hens	3 - 20	375 – 2500
Broilers	3 - 20	375 – 2500
Turkeys	3 - 9	1000 - 1125

Recommended inclusion rates for **Flavo-8** in beef cattle according to feed intake

Recommended dosage: 30 – 50 mg Flavophospholipol head/day

Feed intake per head per day	Flavophospholipol concentration required in the feed (ppm)	Flavo-8 per 1,000 kg of feed
1 kg	40	5000 g
2 kg	20	2500g
3 kg	13.3	1660 g
4 kg	10	1250 g
5 kg	8	1000 g
6 kg	6.7	800 g
7 kg	5.7	700 g
8 kg	5	625 g
9 kg	4.4	550 g
10 kg	4	500 g

BATCH NUMBER:

EXPIRY DATE:

NET WEIGHT: 25 kg

Registration holder: **Animate Animal Health (Pty) Ltd.** (Co.Reg.No.1973/002312/07) 43 Hornbill Avenue, Rooihuiskraal, 0157; PO Box 66698, Highveld, 0169; Phone (+27 12) 661 3485; Fax (+27 12) 661 3486; www.animate.co.za



Animate

Animal Health

VIR DIERLIKE GEBRUIK ALLEEN

FLAVO-8

Reg. Nr. G4151 (Wet 36 van 1947)

VOERVOORMENGSEL

AANDUIDINGS

'n Voerbyvoegsel uitsluitlik vir gebruik as 'n groeistimulant, en vir die verbetering van voeromset by plaasdiere.

Dra by tot die herstel in sensitiviteit van meervoudige medikasie weerstandige *Salmonella typhimurium* en *Escherichia coli* in plaasdiere.

VERSIGTIG

BERGINGSAAWYSINGS

Berg in 'n koel droë plek.

SAMESTELLING

Flavofosfolipol 8 g/kg

WAARSKUWINGS

Hou buite bereik van kinders, ongeligte persone en diere.

Berg weg van voedsel en voer.

Kan vel irritasie veroorsaak.

Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes.

Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

Dit is raadsaam om 'n veiligheidsbril te dra tydens vermenging van die produk. Vermyn inaseming van stof. Moenie eet, drink of rook alvorens hande en gesig deeglik gewas is na hantering van die produk nie.

ONTTREKKINGSPERIODE

Vleis:

Vleisbeeste, varke, skape, bokke en braaikuikens – Nul (0) dae

Melk:

Melk Beeste – Nul (0) dae

Eiers:

Lê-henne – Nul (0) dae

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

Dit is noodsaaklik dat die **FLAVO-8** voormengsel deeglik met die voer gemeng word. Die volgende toedieningsvlakke word vir algemene gebruik aanbeveel:

Gebalanseerde voer	Flavofosfolipol (aktiewe bestanddeel) dpm	FLAVO-8 Voormengsel kg per 1,000 kg voer (8 g aktief per kg)
Melkvervangers	8 - 16	1000 - 2000
Kalwers	8 - 16	1000 - 2000
Volledige bees voer	3 - 20	375 – 2500
Suiwel rantsoene	3 - 20	375 – 2500
Skaap en bok voere	3 - 20	375 – 2500
Klein varkies	8 - 20	1000 – 2500
Alle ander vark voere	3 - 20	375 – 2500
Lê-henne	3 - 20	375 – 2500
Braaikuikens	3 - 20	375 – 2500
Kalkoene	3 - 9	1000 - 1125

Voorgestelde insluiting van **Flavo-8** vir vleisbeeste volgens voer inname.

Aanbevole dosis : 30 – 50 mg Flavofosfolipol/kop/dag

Voeriname per kop per dag	Flavofosfolipol konsentrasie benodig in voer (dpm)	Flavo-8 per 1,000 kg voer
1 kg	40	5000 g
2 kg	20	2500g
3 kg	13.3	1660 g
4 kg	10	1250 g
5 kg	8	1000 g
6 kg	6.7	800 g
7 kg	5.7	700 g
8 kg	5	625 g
9 kg	4.4	550 g
10 kg	4	500 g

LOT NOMMER:

VERVAL DATUM:

NETTO GEWIG: 25 kg

Registrasie houer: Animate Diergesondheid (Edms)Bpk (Co.Reg.Nr.1973/002312/07) 43 Hornbill Laan, Rooihuiskraal, 0157; Posbus 66698, Highveld, 0169; Foon (+27 12) 661 3485; Faks (+27 12) 661 3486; www.animate.co.za

