

Animate

Animal Health

FOR ANIMAL USE ONLY

TM Q®

Reg. No. G3628 (Act 36 of 1947)

FEED PREMIX

INDICATIONS

For the treatment of secondary bacterial infections that complicate primary respiratory infections in poultry, pigs, lambs and calves, caused by bacteria sensitive to oxytetracycline therapy

For the treatment of gastro-enteritis in pigs, lambs and calves caused by bacteria sensitive to oxytetracycline therapy

CAUTION

STORAGE INSTRUCTIONS

Store in a cool dry place.

COMPOSITION

Oxytetracycline 200 g/kg

WARNINGS

All feed medicated with **TM Q®** must be withdrawn 7 days before slaughter.

Keep out of reach of children, uninformed persons and animals.

Store away from food and feed

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder..

WITHDRAWAL PERIOD

Meat: All species - Seven (7) Days

DIRECTIONS FOR USE

Use only as directed.

The dosage rates given below are guidelines only – the inclusion rate should be determined by a veterinarian taking into regard the incidence of disease and the severity of the challenge.

This supplement must be completely dispersed throughout the total feed available to the animal at the recommended rate. The required amount should be mixed with \pm 30 kg of feed, and the resultant be thoroughly blended with the balance of the batch.

Poultry:

Recommended at the following rates:

Therapeutically- 0.5 to 1.0 kg **TM Q®** per ton feed
(i.e. 100 – 200 g oxytetracycline activity per 1,000 kg feed)

Therapy to be continued for 7 – 14 days

Pigs:

Recommended at the following rates:

Therapeutically- 0.5 to 2.0 kg **TM Q®** per ton feed
(i.e. 100 – 400 g oxytetracycline activity per 1,000 kg feed)

Therapy to be continued for 7 – 14 days

Calves & Lambs:

Recommended at the following rates:

Therapeutically- 0.25 to 1.5 kg **TM Q®** per ton feed
(i.e. 50 – 300 g oxytetracycline activity per 1,000 kg feed)

Therapy to be continued for 7 – 14 days

BATCH NUMBER:

EXPIRY DATE:

NET WEIGHT: 25 kg

Registration holder: **Animate Animal Health (Pty)Ltd.** (Co.Reg.No.1973/002312/07)

43 Hornbill Avenue, Rooihuiskraal, 0157; PO Box 66698, Highveld, 0169;

Phone (+27 12) 661 3485; Fax (+27 12) 661 3486; www.animate.co.za



Animate

Animal Health

SLEGS VIR DIERGEBRUIK

TM Q®

Reg. Nr. G3628 (Wet 36 van 1947)

VOER VOORMENGSEL

AANDUIDINGS

Vir die behandeling van sekondêre bakteriese besmetting wat primêre besmettings van die asemhalingskanaal by pluimvee, varke en kalwers kompliseer en veroorsaak word deur bakterieë wat sensitief is vir oksitetrasiklien.

Vir die behandeling van maagdermontsteking by varke en kalwers wat veroorsaak word deur bakterieë wat vir oksitetrasiklien terapie gevoelig is.

VERSIGTIG

BERGINGSAAWYSINGS

Berg in 'n koel droë plek.

SAMESTELLING

Oksitetrasiklien 200 g/kg

WAARSKUWINGS

Alle voere wat met **TM Q®** behandel is, moet 7 dae voor slagtyd onttrek word.

Hou buite bereik van kinders, oningeligte persone en diere.

Berg weg van voedsel en voer.

Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

ONTTREKKINGS PERIODE

Vleis: Alle spesies – Sewe (7) dae

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

Die dosisse hieronder aangedui is slegs riglyne – toediening behoort deur 'n veearts vasgestel te word met inagneming van die siektetoestand en die erns daarvan.

Hierdie byvoegsel moet dwarsdeur die totale beskikbare voer teen die aanbevole peil versprei word. Die nodige hoeveelheid moet eers met ± 30 kg voer gemeng word, en dan moet die mengsel deeglik ingewerk word met die balans van die rantsoen.

Pluimvee:

Aanbeveel teen die volgende peile:

Terapeutiese- 0.5 tot 1.0 kg **TM Q®** per ton voer
(d.w.s. 100 – 200 g oksitetrasiklien
aktiwiteit per 1,000 kg voer)

Terapie moet voortgesit word vir 7 – 14 dae

Varke:

Aanbeveel teen die volgende peile:

Terapeutiese- 0.5 tot 2.0 kg **TM Q®** per ton voer
(d.w.s. 100 – 400 g oksitetrasiklien
aktiwiteit per 1,000 kg voer)

Terapie moet voortgesit word vir 7 – 14 dae

Kalwers & Lammers:

Aanbeveel teen die volgende peile:

Terapeutiese- 0.25 tot 1.5 kg **TM Q®** per ton voer
(d.w.s. 50– 300 g oksitetrasiklien
aktiwiteit per 1,000 kg voer)

Terapie moet voortgesit word vir 7 – 14 dae

LOT NOMMER:

VERVAL DATUM:

NETTO GEWIG: 25 kg

Registrasie houer: Animate Animal Health (Edms)Bpk. (Co.Reg.Nr.1973/002312/07)

43 Hornbill Laan, Rooihuiskraal, 0157; Posbus 66698, Highveld, 0169;

Foon (+27 12) 661 3485; Faks (+27 12) 661 3486; www.animate.co.za

