

Salbutamate® 10%

FOR ANIMAL USE ONLY

SALBUTAMATE® 10%

Reg. No. G4048 (Act 36 of 1947)

FEED PREMIX

INDICATION

CATTLE AND SHEEP: For increased mass gain, increased feed efficiency, improved carcass dressing percentage and improved carcass leanness (lower fat percentage) in cattle and sheep.

CAUTION

STORAGE INSTRUCTIONS

Store in a cool (5°C – 25°C) dry place away from direct sunlight. Keep container well sealed.

COMPOSITION

R-SALBUTAMOL SULPHATE 100 g/kg

WARNINGS

FOR USE IN CATTLE AND SHEEP FINISHER FEEDS ONLY

R-SALBUTAMOL SULPHATE is a cardio active drug and could cause a transient increase in heart rate in persons with coronary artery disease or angina.

Keep out of reach of children, uninformed persons and animals.

Store away from food, drink and other animal feeds.

Keep container in a well-ventilated area.

When mixing and handling the product, use protective clothing, impervious gloves and a dust mask. Operators should wash their hands thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Not to be used after expiry date indicated on the bag. Medicated feed must be used within 3 months after mixing.

WITHDRAWAL PERIOD

Cattle - Meat: Two (2) Days

Sheep – Meat: One (1) Day

Milk: Not for use in animals producing milk for human consumption

DIRECTIONS FOR USE

Use only as directed.

This supplement must be completely dispersed throughout the total feed available to the animal at the recommended rate. The required amount should be mixed with ± 30 kg of feed, and the resultant 30kg be thoroughly blended with the balance of the final batch.

DOSAGE

Cattle:

Include **Salbutamate® 10%** at 75 to 150 mg R-Salbutamol sulphate per animal per day in the final feed for the last 30 to 40 days of the finishing phase.

Sheep:

Include **Salbutamate 10%** at 100 to 150g/ton of feed in the final feed for the last 28 days of the finishing phase.

SLEGS VIR DIEREGBEUK

SALBUTAMATE® 10%

Reg. Nr. G4048 (Wet 36 van 1947)

VOER VOORMENGSEL

INDIKASIE

BEESTE EN SKAPE: Vir verhoogde massa toename, verbeterde voeromset, verbeterde karkas uitslagpersentasie en verbeterde karkas maerheid (laer vet persentasie) in beeste en skape.

VERSIGTIG

BERGINGS INSTRUKSIES

Stoor in 'n koel (5°C – 25°C) droë plek weg van direkte sonlig. Hou verpakking goed geseël.

SAMESTELLING

R-SALBUTAMOL SULFAAT 100 g/kg

WAARSKUWING

SLEGS VIR GEBRUIK IN BEES- EN SKAAPAFRONDINGSVOER

R-SALBUTAMOL SULFAAT is 'n kardio-aktiewe middel en kan 'n verhoogde harttempo tot gevolg hê by mense met korinêre hartsiektes of angina.

Hou buite die bereik van kinders, oningeligte persone en diere.

Berg weg van voedsel, drank en ander dierevoere.

Berg houer in 'n goed geventileerde plek.

Gebruik beskermende klere, handskoene en 'n stofmasker wanneer produk gemeng en hanteer word. Operateurs moet hulle hande deeglik met seep en water was na hantering van die produk. As produk per ongeluk met oë in kontak kom moet dit onmiddellik deeglik met water uitgespoel word.

Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit misluk as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORG MAATREËLS

Moet nie na verval datum, soos aangedui op die verpakking, gebruik word nie. Gemedikeerde voer moet binne 3 maande van vermenging gebruik word.

ONTTREKKINGSPERIODE

Bees – Vleis: Twee (2) Dae

Skaap – Vleis: Een (1) Dag

Melk: Nie vir die gebruik in diere wat melk produseer vir menslike gebruik nie

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui.

Die produk moet deeglik gemeng word met die totale voer wat beskikbaar gestel word aan die dier teen die aanbevole vlakke.

Die korrekte hoeveelheid produk moet met ± 30 kg voer vermeng word en die 30kg mengsel moet dan weer deeglik in die res van die finale mengsel ingemeng word.

DOSERING

Bees:

Meng **Salbutamate® 10%** in die finale voer teen 75 tot 150 mg R-Salbutamol sulfaat per dier per dag vir die laaste 30 – 40 dae van die afrondingsperiode.

Skaap:

Meng **Salbutamate® 10%** in die finale voer teen 110 tot 150 g per ton voer vir die laaste 28 dae van die afrondingsperiode.



Registration holder/Registration holder:

Animate Animal Health (Pty) Ltd. (Co.Reg.No.1973/002312/07)
43 Hornbill Avenue, Rooihuiskraal, 0157; PO Box 66698, Highveld, 0169;
Phone (+27 12) 661 3485; Fax (+27 12) 661 3486; www.animate.co.za

Animate
Animal Health

